



## The New Integral Transform” Iman Transform” of Bessel’s Function with Application for Evaluating Definite Integrals

Huda M. Khalat

Mathematics Department, Faculty of Sciences, University- Sabratha /Libya  
[huda.khalat@sabu.edu.ly](mailto:huda.khalat@sabu.edu.ly)

### ARTICLE INFO

#### Article history:

Received: 12/02/2025

Received in revised form 21/04/2025

Accepted: 11/05/2025

### ABSTRACT

In the advance time, Bessel’s function play role in solving many problems of engineering and scientific challenges. Including equations like the wave equation, heat equation, Laplace equation, Helmholtz equation, and Schrodinger equation in spherical or cylindrical coordinates. In this paper, author present Iman transform of Bessel’s functions with application for evaluating definite integrals, and exploring its applications and providing examples of problems involving Bessel’s functions for integral calculation, thereby minimizing computational efforts.

**Keywords:** Bessel’s function, Iman transform, Convolution theorem, Inverse Iman transform, Definite integral.

### 1. Introduction

Recently, integral transforms such as the Laplace transform [1,2], Elzaki transform [3,4], Kamal transform [5-7], Mahgoub transform [8], Shehu transform [9], Sawi transform [10], SEE Transform [11], Abaoub Skhem transform [12,13] etc, have played a crucial role in solving advanced problems in fields such as mathematics, physics, chemistry, social science,

astronomy, biology, nuclear science, and engineering. These transforms are particularly useful for addressing complex challenges like differential equations, integro-differential equations, and population dynamics. Similarly, the widespread use of integral transform methods continues to prove valuable in

solving various engineering and scientific problems [14].

Researchers have successfully applied these methods to ordinary differential equations [15], partial differential equation [16,17], Volterra integral equations [18], and Volterra integro-differential equations [19]. For instance, Priyyanka and Aggarwal [20] utilized the Risi transforms to model bacterial growth, while Aggarwal and colleagues [21] have explored several transforms of Bessel functions. Building on this foundation. Application of Anuj transform for handling linear Volterra integro-differential equations and first order consecutive chemical reaction was given by [22,23]. Khalat and Ali [24] gave the solution of Elliptical Curves equation.

Bessel's functions are widely applied in solving problems across various fields, including mathematical physics, acoustics, engineering, and sciences like heat transfer, fluid mechanics, vibrations, stress analysis, hydrodynamics, and flux distribution in nuclear reactors, among other etc.

The purpose of this study is to determine the Iman transform of Bessel functions of the first kind of orders zero, one, and two and to demonstrate the advantages of using the Iman transform for evaluating integrals that involve Bessel's function.

## 2. Definition and Standard Results

### 2.1. Iman transform

**Definition:** If  $f(t) \in F, t \geq 0$  then Iman transform of  $f(t)$  is defined as [25]

$$I\{f(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(t) dt = K(v), v > 0 \quad (1)$$

Where  $I$  is called the Iman transform operator.

### 2.2. properties of Iman transform

#### 2.2. a) Change of scale

If  $I\{f(t)\} = K(v)$  then  $I\{f(at)\} = aK(av)$ , where  $a$  is arbitrary constant.

**proof:** By the definition of Iman transform, we have

$$I\{f(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(t) dt, \text{ then}$$

$$I\{f(at)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(at) dt$$

Put  $at = p \rightarrow adt = dp$  in above equation, we have

$$I\{f(at)\} = \frac{1}{av^2} \int_0^\infty e^{-v^2 \frac{p}{a}} f(p) dp$$

$$I\{f(at)\} = \left[ \frac{1}{a^2 \left(\frac{v}{\sqrt{a}}\right)^2} \int_0^\infty e^{-\left(\frac{v}{\sqrt{a}}\right)^2 p} f(p) dp \right]$$

$$= \frac{1}{a^2} K\left(\frac{v}{\sqrt{a}}\right)$$

thus if  $I\{f(t)\} = K(v)$ , then

$$I\{f(at)\} = \frac{1}{a^2} K\left(\frac{v}{\sqrt{a}}\right).$$

#### 2.2. b) Linearity Property

If  $I\{f(t)\} = K(v)$  and  $I\{g(t)\} = H(v)$

We have  $I\{af(t) + bg(t)\} = aI\{f(t)\} + bI\{g(t)\}$

Then  $I\{af(t) + bg(t)\} = aK(v) + bH(v)$

Where  $a, b$  are arbitrary constants.

**Proof:** By the definition of Iman transform, we have

$$I\{f(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(t) dt$$

$$I\{af(t) + bg(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} [af(t) + bg(t)] dt$$

$$= a \left[ \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(t) dt \right] + b \left[ \frac{1}{v^2} \int_0^\infty e^{-v^2 t} g(t) dt \right]$$

Then  $I\{af(t) + bg(t)\} = aI\{f(t)\} + bI\{g(t)\}$

And  $I\{af(t) + bg(t)\} = aK(v) + bH(v)$

Where  $a, b$  are arbitrary constants.

### 3. Iman transform of the derivatives of the function $f(t)$ [25-27]

Let  $K(v)$  be Iman transform of  $[I\{f(t)\} = K(v)]$ , then

$$\begin{aligned}
(a) \quad I \left\{ \frac{df(t)}{dt} \right\} &= v^2 K(v) - \frac{1}{v^2} f(0) \\
(b) \quad I \left\{ \frac{d^2 f(t)}{dt^2} \right\} &= v^4 K(v) - f(0) - \frac{1}{v^2} f'(0) \\
(c) \quad I \left\{ \frac{d^n f(t)}{dt^n} \right\} &= v^{2n} K(v) - \\
&\quad \sum_{k=0}^{n-1} \frac{1}{v^{4-2n+2k}} f^{(k)}(0).
\end{aligned}$$

#### 4. Convolution of two functions

The convolution of two functions  $f(t)$  and  $g(t)$  is represented as  $f(t) * g(t)$

and is defined by:

$$\begin{aligned}
f(t) * g(t) &= f * g = \int_0^t f(t)g(x-t)dt \\
&= \int_0^t g(t)f(x-t)dt
\end{aligned}$$

##### 4.1. Convolution theorem for Iman transforms

If  $I\{f(t)\} = K_1(v)$  and  $I\{g(t)\} = K_2(v)$ , then  $I\{f(t) * g(t)\} = v^2 I\{f(t)\} * I\{g(t)\} = v^2 K_1(v) \cdot K_2(v)$ .

**Proof:** By the definition of Iman transform, we have

$$I\{f(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} f(t) dt, \text{ then}$$

$$I\{f(t) * g(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} [f(t) * g(t)] dt$$

From the definition of convolution of two functions, we obtain

$$I\{f(t) * g(t)\} = \frac{1}{v^2} \int_0^\infty e^{-v^2 t} \left[ \int_0^t f(x)g(t-x) \right] dt$$

Changing the order of integration gives

$$I\{f(t) * g(t)\} = \frac{1}{v^2} \int_0^\infty f(x) \left[ \int_x^\infty e^{-v^2 t} g(t-x) dt \right] dx$$

Put  $t-x = u$  so that  $dt = du$  in above equation, we have

$$\begin{aligned}
I\{f(t) * g(t)\} &= \frac{1}{v^2} \int_0^\infty f(x) \left[ \int_x^\infty e^{-v^2(u+x)} g(u) du \right] dx \\
I\{f(t) * g(t)\} &= \frac{1}{v^2} \int_0^\infty f(x) e^{-v^2 x} \left[ \int_0^\infty e^{-v^2 u} g(u) du \right] dx
\end{aligned}$$

$$\begin{aligned}
I\{f(t) * g(t)\} &= \int_0^\infty f(x) e^{-v^2 x} [I\{g(t)\}] dx \\
I\{f(t) * g(t)\} &= v^2 I\{g(t)\} \left[ \frac{1}{v^2} \int_0^\infty e^{-v^2 x} f(x) dx \right] \\
I\{f(t) * g(t)\} &= v^2 I\{f(t)\} * I\{g(t)\} = \\
&= v^2 K_1(v) \cdot K_2(v).
\end{aligned}$$

Table .1. Iman transform of some elementary functions [25]

| S.N | $f(t)$         | $I\{f(t)\} = K(v)$             |
|-----|----------------|--------------------------------|
| 1.  | 1              | $\frac{1}{v^4}$                |
| 2.  | t              | $\frac{1}{v^6}$                |
| 3.  | $t^n, n \in N$ | $\frac{n!}{v^{2n+4}}$          |
| 4.  | $t^n, n > -1$  | $\frac{\Gamma(n+1)}{v^{2n+4}}$ |
| 5.  | $e^{at}$       | $\frac{1}{v^4 - av^2}$         |
| 6.  | $e^{-at}$      | $\frac{1}{v^4 + av^2}$         |
| 7.  | $\sin at$      | $\frac{a}{v^2(v^4 + a^2)}$     |
| 8.  | $\cos at$      | $\frac{1}{v^4 + a^2}$          |
| 9.  | $\sinh at$     | $\frac{a}{v^2(v^4 - a^2)}$     |
| 10. | $\cosh at$     | $\frac{1}{v^4 - a^2}$          |

#### 5. Inverse Iman transform

**Definition:** If  $I\{f(t)\} = K(v)$  then  $f(t)$  is called the inverse Iman transform of  $K(v)$  and mathematically it is denoted by

$$f(t) = I^{-1}\{K(v)\}$$

Where the operator  $I^{-1}$  is the inverse Iman transform.

Table .2. Inverse Iman transform of some elementary functions [25-28]

| S.N | $K(v)$                        | $f(t) = I^{-1}\{K(v)\}$   |
|-----|-------------------------------|---------------------------|
| 1.  | $\frac{1}{v^4}$               | 1                         |
| 2.  | $\frac{1}{v^6}$               | t                         |
| 3.  | $\frac{1}{v^{2n+4}}, n \in N$ | $n! t^n$                  |
| 4.  | $\frac{1}{v^{2n+4}}, n > -1$  | $\Gamma(n+1) t^n, n > -1$ |
| 5.  | $\frac{1}{v^4 - av^2}$        | $e^{at}$                  |
| 6.  | $\frac{1}{v^4 + av^2}$        | $e^{-at}$                 |

|     |                            |                      |
|-----|----------------------------|----------------------|
| 7.  | $\frac{1}{v^2(v^4 + a^2)}$ | $\frac{\sin at}{a}$  |
| 8.  | $\frac{1}{v^4 + a^2}$      | $\cos(at)$           |
| 9.  | $\frac{1}{v^2(v^4 - a^2)}$ | $\frac{\sinh at}{a}$ |
| 10. | $\frac{1}{v^4 - a^2}$      | $\cosh at$           |

## 6. Bessel's functions of first kind of different order [29,30]

Bessel's function of order  $n$ , where  $n \in \mathbb{N}$  is given by

$$J_n(t) = \frac{t^n}{2^n n!} \left[ 1 - \frac{t^2}{2 \cdot (2n+2)} + \frac{t^4}{2 \cdot 4 \cdot (2n+2)(2n+4)} - \frac{t^6}{2 \cdot 4 \cdot 6 \cdot (2n+2)(2n+4)(2n+6)} + \dots \right] \quad (2)$$

When  $n = 0$ , we have Bessel's function of zero order and it denoted by  $J_0(t)$  and it is given by

$$J_0(t) = 1 - \frac{t^2}{2^2} + \frac{t^4}{2^2 \cdot 4^2} - \frac{t^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \quad (3)$$

And when  $n = 1$ , we have Bessel's function of order one, it is given by

$$J_1(t) = \frac{t}{2} - \frac{t^3}{2^3 \cdot 4} + \frac{t^5}{2^2 \cdot 4^2 \cdot 6} - \frac{t^7}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8} + \dots \quad (4)$$

Another form of Equation (3) can be written as

$$J_1(t) = \frac{t}{2} - \frac{t^3}{2^3 \cdot 2!} + \frac{t^5}{2^5 \cdot 2! \cdot 3!} - \frac{t^7}{2^7 \cdot 3! \cdot 4!} + \dots \quad (5)$$

For  $n = 2$ , Bessel's function of order two, it is given by

$$J_2(t) = \frac{t^2}{2 \cdot 4} - \frac{t^4}{2^2 \cdot 4 \cdot 6} + \frac{t^6}{2^2 \cdot 4^2 \cdot 6 \cdot 8} - \frac{t^8}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8 \cdot 10} + \dots \quad (6)$$

## 7. Relation between $J_0(t)$ and $J_1(t)$ [29]

$$\frac{d}{dt} [J_0(t)] = -J_1(t) \quad (7)$$

## 8. Relation between $J_0(t)$ and $J_2(t)$ [30]

$$J_2(t) = J_0(t) + 2J_0''(t) \quad (8)$$

## 9. Iman transform of Bessel's functions of first kind

### 9.1. Iman transform of $J_0(t)$

Applying Iman transform of equation (3), both sides, we have:

$$\begin{aligned} I[J_0(t)] &= I[1] - \frac{1}{2^2} I[t^2] + \frac{1}{2^2 \cdot 4^2} I[t^4] \\ &\quad - \frac{1}{2^2 \cdot 4^2 \cdot 6^2} I[t^6] + \dots \\ &= \frac{1}{v^4} - \frac{1}{2^2} \cdot \frac{1}{v^8} + \frac{1}{2^2 \cdot 4^2} \cdot \frac{1}{v^{12}} - \frac{1}{2^2 \cdot 4^2 \cdot 6^2} \cdot \frac{1}{v^{16}} + \dots \\ &= \frac{1}{v^4} \left[ 1 - \frac{1}{2^2} \cdot \frac{2!}{v^4} + \frac{1}{2^2 \cdot 4^2} \cdot \frac{4!}{v^8} - \frac{1}{2^2 \cdot 4^2 \cdot 6^2} \cdot \frac{6!}{v^{12}} + \dots \right] \\ &= \frac{1}{v^4} \left[ 1 - \frac{1}{2} \left( \frac{1}{v^4} \right) + \frac{3 \cdot 1}{2 \cdot 4} \left( \frac{1}{v^4} \right)^2 - \frac{5 \cdot 3 \cdot 1}{2 \cdot 4} \left( \frac{1}{v^4} \right)^3 + \dots \right] \\ &= \frac{1}{v^4} \left( 1 + \frac{1}{v^4} \right)^{-\frac{1}{2}} \\ &= \frac{1}{v^2 \sqrt{1 + v^4}} \quad (9) \end{aligned}$$

### 9.2. Iman transform of $J_1(t)$

From (7), we have  $\frac{d}{dt} [J_0] = -J_1(t)$

Then  $I[J_1(t)] = -I \left[ \frac{d}{dt} J_0(t) \right]$

$$\begin{aligned} &= - \left[ \frac{-1}{v^2} J_0(0) + v^2 I[J_0(t)] \right] \\ &= \frac{1}{v^2} J_0(0) - v^2 I[J_0(t)] \\ &= \frac{1}{v^2} (1) - \frac{v^2}{v^4 \sqrt{1 + \frac{1}{v^4}}} \\ &= \frac{1}{v^2} \left[ 1 - \frac{v^2}{\sqrt{1 + v^4}} \right] \quad (10) \end{aligned}$$

### 9.3. Iman transform of $J_2(t)$

Taking Iman transform of equation (8), both sides, we get

$$\begin{aligned} I[J_2(t)] &= I[J_0(t)] + 2I[J_0''(t)] \\ &= \frac{1}{v^2\sqrt{1+v^4}} + 2\left[v^4 I[J_0(t)] - \frac{1}{v^2} J_0'(0) - J_0(0)\right] \\ &= \frac{1}{v^2\sqrt{1+v^4}} + 2\left[\frac{v^2}{v^2\sqrt{1+v^4}} - 0 - 1\right] \\ &= \frac{1}{v^2\sqrt{1+v^4}} + \frac{2}{\sqrt{1+v^4}} - 2 \\ &= \frac{1+2v^2-2v^2\sqrt{1+v^4}}{v^2\sqrt{1+v^4}} \end{aligned} \quad (11)$$

### 9.4. Iman transform of $J_0(at)$

$$\text{Since } I[J_0(t)] = \frac{1}{v^2\sqrt{1+v^4}}$$

Now, using change of scale property of Iman transform, we have:

$$\begin{aligned} I[J_0(at)] &= \frac{1}{a^2} \left[ \frac{1}{\left(\frac{v}{\sqrt{a}}\right)^2 \sqrt{1+\left(\frac{v}{\sqrt{a}}\right)^4}} \right] \\ &= \frac{1}{v^2} \left[ \frac{1}{\sqrt{a^2+v^4}} \right] \end{aligned} \quad (12)$$

### 9.5. Iman transform of $J_1(at)$

$$\text{Since } I[J_1(t)] = \frac{1}{v^2} \left[ 1 - \frac{v^2}{\sqrt{1+v^4}} \right]$$

$$\begin{aligned} \text{Then } I[J_2(t)] &= \frac{1}{a^2} \left[ \frac{1}{\left(\frac{v}{\sqrt{a}}\right)^2} \left[ 1 - \frac{\left(\frac{v}{\sqrt{a}}\right)^2}{\sqrt{1+\left(\frac{v}{\sqrt{a}}\right)^4}} \right] \right] \\ &= \left[ \frac{1}{av^2} - \frac{1}{a^3\sqrt{a^2+v^4}} \right] \end{aligned} \quad (13)$$

### 9.6. Iman transform of $J_2(at)$

$$\text{Since } I[J_2(t)] = \frac{1+2v^2-2v^2\sqrt{1+v^4}}{v^2\sqrt{1+v^4}}$$

$$\begin{aligned} \text{Then } I[J_2(at)] &= \frac{1}{a^2} \left[ \frac{1+2\frac{v^2}{a^2}-2\frac{v^2}{a^2}\sqrt{a^2+v^4}}{\frac{v^2}{a^2}\sqrt{a^2+v^4}} \right] \\ &= \frac{1}{a^2v^2} \left[ \frac{a^2+2av^2-2v^2\sqrt{a^2+v^4}}{\sqrt{a^2+v^4}} \right] \end{aligned} \quad (14)$$

## 10. First shifting Theorem for Iman transform

Let  $I[f(t)] = k(v)$  and  $a \in R$ , then

$$I[e^{at}f(t)] = \left(1 - \frac{a}{v^2}\right) k\left(v\sqrt{1 - \frac{a}{v^2}}\right)$$

**Proof:** From definition of a new integral transform, we have

$$\begin{aligned} I[e^{at}f(t)] &= \frac{1}{v^2} \int_0^\infty e^{at} f(t) e^{-v^2t} dt \\ &= \frac{1}{v^2} \int_0^\infty f(t) e^{-[v^2-a]t} dt \\ &= \left(1 - \frac{a}{v^2}\right) \frac{1}{\left(v\sqrt{1 - \frac{a}{v^2}}\right)^2} \int_0^\infty f(t) e^{-\left(v\sqrt{1 - \frac{a}{v^2}}\right)^2 t} dt \\ &= \left(1 - \frac{a}{v^2}\right) k\left(v\sqrt{1 - \frac{a}{v^2}}\right) \end{aligned} \quad (15)$$

### 10.1. Iman transform of $e^{at}J_0(t)$

From  $J_0(t)$  we have

$$I[J_0(t)] = \frac{1}{v^2\sqrt{1+v^4}}$$

Using first shifting property we get

$$\begin{aligned} I[e^{at}J_0(t)] &= \left(1 - \frac{a}{v^2}\right) \left[ \frac{1}{v^2\left(1 - \frac{a}{v^2}\right)\sqrt{1+v^4\left(1 - \frac{a}{v^2}\right)^2}} \right] \\ &= \left[ \frac{1}{v^2\sqrt{1+v^4-2av^2+a^2}} \right] \end{aligned} \quad (16)$$

### 10.2. Iman transform of $e^{at}J_1(t)$

Since  $I[J_1(t)] = \frac{1}{v^2} \left[ 1 - \frac{v^2}{\sqrt{1+v^4}} \right]$

Thus  $I[e^{at}J_1(t)] =$

$$\left(1 - \frac{a}{v^2}\right) \left( \frac{1}{v^2 \left(1 - \frac{a}{v^2}\right)} \right) \left[ 1 - \frac{v^2 \left(1 - \frac{a}{v^2}\right)}{\sqrt{1+v^4 \left(1 - \frac{a}{v^2}\right)^2}} \right]$$

$$= \frac{1}{v^2} \left[ \frac{\sqrt{1 + (v^2 - a)^2} - v^2 + a}{\sqrt{1 + (v^2 - a)^2}} \right] \quad (17)$$

### 10.3. Iman transform of $e^{at}J_2(t)$

Since  $I[J_2(t)] = \frac{1+2v^2-2v^2\sqrt{1+v^4}}{v^2\sqrt{1+v^4}}$

Then  $I[e^{at}J_2(t)] =$

$$\left(1 - \frac{a}{v^2}\right) \left[ \frac{1+2v^2\left(1 - \frac{a}{v^2}\right) - 2v^2\left(1 - \frac{a}{v^2}\right)\sqrt{1+v^4\left(1 - \frac{a}{v^2}\right)^2}}{\left(1 - \frac{a}{v^2}\right)\sqrt{1+v^4\left(1 - \frac{a}{v^2}\right)^2}} \right]$$

$$= \frac{1 + 2(v^2 - a) - 2(v^2 - a)\sqrt{1 + (v^2 - a)^2}}{\sqrt{1 + (v^2 - a)^2}} \quad (18)$$

## 11. Numerical applications

In this section, some examples are provided to illustrate the effectiveness of Iman transform of Bessel's functions in evaluating the integral that involve Bessel's functions.

**Application 1.** Evaluate the integral

$$E(t) = \int_0^t J_0(u)J_0(t-u)du \quad (19)$$

Taking Iman transform to both sides of equation (19), we have

$$I\{E(t)\} = I\left\{ \int_0^t J_0(u)J_0(t-u)du \right\} \quad (20)$$

Applying convolution theorem of Iman transform on (20), we have

$$I\{E(t)\} = v^2 I\{J_0(t)\} I\{J_0(t)\}$$

$$= v^2 \left[ \frac{1}{v^2 \sqrt{1+v^4}} \right] \left[ \frac{1}{v^2 \sqrt{1+v^4}} \right]$$

$$= \frac{1}{v^2(1+v^4)} \quad (21)$$

Now operating inverse Iman transform on both side on (21), we have

$$E(t) = I^{-1} \left\{ \frac{1}{v^2(1+v^4)} \right\} = \sin(t) \quad (22)$$

Which is the exact solution needed for (19).

**Application 2.** Evaluate the integral

$$E(t) = \int_0^t J_0(u)J_1(t-u)du \quad (23)$$

Taking Iman transform to both sides of equation (23), we have

$$I\{E(t)\}$$

$$= I \left\{ \int_0^t J_0(u)J_1(t-u)du \right\} \quad (24)$$

Using convolution theorem of Iman transform on (24), we have

$$I\{E(t)\} = v^2 I\{J_0(t)\} I\{J_1(t)\}$$

$$= v^2 \left[ \frac{1}{v^2 \sqrt{1+v^4}} \right] \left[ \frac{1}{v^2} \left[ 1 - \frac{v^2}{\sqrt{1+v^4}} \right] \right]$$

$$= \frac{1}{v^2 \sqrt{1+v^4}} - \frac{1}{1+v^4} \quad (25)$$

Take inverse to both sides on (25), we get

$$E(t) = I^{-1} \left\{ \frac{1}{v^2 \sqrt{1+v^4}} \right\} - I^{-1} \left\{ \frac{1}{1+v^4} \right\}$$

$$= J_0(t) - \cos(t) \quad (26)$$

The exact solution of (23) that is required.

**Example 3.** Evaluate the integral

$$E(t) = \int_0^t J_1(t-u)du \quad (27)$$

Applying the Iman transform on both sides of equation (27), we have

$$I\{E(t)\} = I\left\{\int_0^t J_1(t-u)du\right\} \quad (28)$$

Using convolution theorem of Iman on (28), we have

$$\begin{aligned} I\{E(t)\} &= v^2 I\{1\} I\{J_1(t)\} \\ &= v^2 \left[ \frac{1}{v^4} \right] \left[ \frac{1}{v^2} \left[ 1 - \frac{v^2}{\sqrt{1+v^4}} \right] \right] \\ &= \frac{1}{v^4} - \frac{1}{v^2 \sqrt{1+v^4}} \end{aligned} \quad (29)$$

Now taking inverse Iman transform on both sides on (29), we have

$$\begin{aligned} E(t) &= I^{-1}\left\{\frac{1}{v^4}\right\} - I^{-1}\left\{\frac{1}{v^2 \sqrt{1+v^4}}\right\} \\ &= 1 - J_0(t) \end{aligned} \quad (30)$$

Which is the required solution of equation (27).

## 12. Conclusion

In this work, the Iman transform is employed to solve Bessel functions of the first kind of order zero, one, and two, denoted as  $J_0(t)$ ,  $J_1(t)$ ,  $J_2(t)$ ,  $J_0(at)$ ,  $J_1(at)$  and  $J_2(at)$ . Additionally, they obtained the Iman transform for expressions involving these functions, such as  $e^{kt}J_0(t)$ ,  $e^{kt}J_1(t)$ ,  $e^{kt}J_2(t)$ , using the translation and scaling properties of the Iman transform. These findings are significant for evaluating improper integrals that include Bessel functions in the integrand. Additionally, we explore several properties and theorems related to the Iman transform, we applied this transform to three different examples. The results of this paper may also be useful in future research for solving Bessel's equations.

## 13. References

1. Aggarwal, S., & Chaudhary, R. (2019). A comparative study of Mohand and Laplace transforms. *Journal of Emerging*

*Technologies and Innovative Research*, 6(2), 230-240.

2. Frasser, C.E., Laplace Transform and Systems of Ordinary Differential Equations. Jan2019, 1-34.
3. Aggarwal, S., Gupta, A. R., & Kumar, A. (2019). Elzaki transform of error function. *Global Journal of Engineering Science and Researches*, 6(5), 412-422.
4. Aggarwal, S., Bhatnagar, K., & Dua, A. (2019). Dualities between Elzaki transform and some useful integral transforms. *International Journal of Innovative Technology and Exploring Engineering*, 8(12), 4312-4318.
5. Aggarwal, S., Chauhan, R., & Sharma, N. (2018). A new application of Kamal transforms for solving linear Volterra integral equations. *International Journal of Latest Technology in Engineering, Management & Applied Science*, 7(4), 138-140.
6. Aggarwal, S., Sharma, N., & Chauhan, R. (2020). Duality relations of Kamal transform with Laplace, Laplace–Carson, Aboodh, Sumudu, Elzaki, Mohand and Sawi transforms. *SN Applied Sciences*, 2(1), 135.
7. Aggarwal, S., Kamal transforms of Bessel's functions, *International Journal of Research and Innovation in Applied Science*, 3(7), 1-4, 2018.
8. Aggarwal, S., Sharma, N., & Chauhan, R. (2018). Solution of linear Volterra integro-differential equations of second kind using Mahgoub transform. *International Journal of Latest Technology in Engineering*,

- Management & Applied Science, 7(5), 173-176.
9. Aggarwal, S., Vyas, A. and Sharma, S.D., Primitive of first kind Volterra integro-differential equation using Shehu transform, International Journal of Research and Innovation in Applied Science, 5(8), 31-38, 2020.
  10. Aggarwal, S., Sharma, S.D. and Vyas, A., Sawi transform of Bessel's functions with application for evaluating definite integrals, International Journal of Latest Technology in Engineering, Management & Applied Science, 9(7), 12-18, 2020.
  11. Mansour, E. A., Mehdi, S. A., kuffi, E. A. kuffi, "The New Integral Transform "SEE Transform" of Bessel's Functions Journal of Computer and Mathematics Education, Vol.12 No.13 (2021), 3898-3905.
  12. Khalat H.M., Mubayrash, A. O., A New Cryptographic Scheme Using Integral Transform and Matrix Decomposition, The Libyan Journal of Science-University of Tripoli, Vol.27, No.02 (2024), 56-64.
  13. Mubayrash, A. O., Khalat, H.M., New method for Cryptography using Abaoub-Shkheam Transform, The Libyan Journal of Science-University of Tripoli, 25, (02), (2022), 35-39.
  14. S. Maitama and W. Zhao, New integral transform: Shehu transform a generalization of Sumudu and Laplace transform for solving differential equations, International Journal of Analysis and Applications, 17(2), 167-190, 2019.
  15. Aggarwal, S., Sharma, N., Chauhan, R., Gupta, A.R. and Khandelwal, A., A new application of Mahgoub transforms for solving linear ordinary differential equations with variable coefficients, Journal of computer and Mathematical Sciences, 9(6), 520-525, 2018.
  16. Ahmed, S.A., Elzaki, T.M., Elbadri, M. and Mohamed, M.Z., Solution of partial differential equations by new double integral transform (Laplace-Sumudu transform), Ain Shams Engineering Journal, 12(2021), 4045-4049. <https://doi.org/10.1016/j.asej.2021.02.032>
  17. Chauhan, R., & Aggarwal, S. (2018). Solution of linear partial integro-differential equations using Mahgoub transform. Periodic Research, 7(1), 28-31.
  18. Chauhan, R. and Aggarwal, S., Laplace transform for convolution type linear Volterra integral equation of second kind, Journal of Advanced Research in Applied Mathematics and Statistics, 4(3&4), 1-7, 2019.
  19. Gupta, A. R., Aggarwal, S., & Agrawal, D. (2018). Solution of linear partial integro-differential equations using Kamal transform. International Journal of Latest Technology in Engineering, Management & Applied Science, 7(7), 88-91.
  20. Priyanka and Aggarwal, S., Solution of the model of the bacteria growth via Rishi transform, Journal of Advanced Research in Applied Mathematics and

- Statistics, 7(1&2), 5-11, 2022.  
<https://doi.org/10.24321/2455.7021.202202>
21. Kumar, A., Bansal, S. and Aggarwal, S., Determination of the blood glucose concentration of a patient during continuous intravenous injection using Anuj transform, Neuroquantology, 19(12), 303-306, 202
  22. Kumar, A., Bansal, S. and Aggarwal, S., A new novel integral transform “Anuj transform” with application, Design Engineering, 2021(9), 12741-12751, 2021.
  23. Kumar, A., Bansal, S. and Aggarwal, S., Determination of the concentrations of the reactants of first order consecutive chemical reaction using Anuj transform, European Chemical Bulletin, 12(7), 2528-2534, 2023.
  24. Khalat, H. M., Ali, N. F, Application of RSA Cryptography and Elliptical Curves in Public Key Cryptography, Journal of Faculties of Education, the Twenty-Five Issue March (2022),1-20.
  25. Patil, D. P., Bagul, A. D., Sonawane, V. S., Iman Transform for the Solution in Models in Health Sciences, Journal of Emerging Technologies and innovative Research, Vol.10 Issue12, (12), (2023), a662-a669.
  26. Almardy, I.A., The New Integral Transform “Iman Transform”, Journal of Advanced Research in science, Vol.4,(03), 2023,1-5.
  27. On the Iman Transform and Systems of Ordinary Differential Equations I. A. Almardy ,R. A. Farah, M. A. Elkeer, Volume 3, Issue 1, February 2023, International Journal of Advanced Research in Science, Communication and Technology (IJARSCT), ISSN (Online) 2581-9429, Copyright to IJARSCT DOI: [10.48175/568580](https://doi.org/10.48175/568580) [www.ijarsct.co.in](http://www.ijarsct.co.in).
  28. I. A. Almardy, S. Y. Eltayeb, S. H. Mohamed, M. A. Alkerr, A. K. Osman, H. A. Albushra, A Comparative Study of Iman and Laplace Transforms to Solve Ordinary Differential Equations of First and Second Order, ISSN 1, Volume 3, (2023), pp323-328. International Journal of Advanced Research in Science, Communication and Technology (IJARSCT).
  29. Aggarwal, S., N. and Chauhan, R., Mahgoub transform of Bessel’s functions, International Journal of Latest Technology in Engineering, Management &Applied Science,7(8),32-36,2018.
  30. Aggarwal, S., Elzaki transform of Bessel’s functions, Global Journal of Engineering Science and Researches,5(8),45-51,2018.